

考 試 科 目	基礎數學	系 所 別	統計學系	考 試 時 間	2 月 9 日 (三) 第一節
---------	------	-------	------	---------	-----------------

1. (36pts). For each of the following statements, determine whether it is true (○) or false (×). (True or false questions. Do not give explanation.)

- (a) $\lim_{x \rightarrow 0} \frac{|3x-1|-|3x+1|}{x}$ does not exist.
- (b) If functions $f(x)$ and $g(x)$ are continuous on $[a, b]$, then the following definite integrals exist and $\int_a^b f(x)g(x)dx = \int_a^b f(x)dx \int_a^b g(x)dx$.
- (c) The equations (i) $x^2 + y^2 = 9$, (ii) $x = -3 \cos 2t, y = -3 \sin 2t$ ($0 \leq t \leq \pi$), and (iii) $r = 3$ all have the same graph.
- (d) Suppose that the radius of convergence of the power series $\sum_{k=0}^{\infty} a_k x^k$ is R_0 for $0 < R_0 < \infty$. Then the radius of convergence of the power series $\sum_{k=0}^{\infty} a_k x^{2k}$ is R_0^2 .
- (e) The function $f(x, y) = \begin{cases} \frac{x^2 y}{x^2 + y^4} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$ is continuous at $(0, 0)$.
- (f) Suppose $f(x, y) : \mathcal{R} \times \mathcal{R} \rightarrow \mathcal{R}$ has continuous second partial derivatives on $(-\delta, \delta) \times (-\delta, \delta)$ for some $\delta > 0$, and suppose that $f_x(0, 0) = 0$ and $f_y(0, 0) = 0$. If the matrix $\begin{bmatrix} f_{xx}(0, 0) & f_{xy}(0, 0) \\ f_{yx}(0, 0) & f_{yy}(0, 0) \end{bmatrix}$ is positively definite, then f has a local minimum at $(0, 0)$.
- (g) A homogeneous system of four equations in six unknowns always has nontrivial solutions.
- (h) If A is nonsingular and $A^{-1} = A^T$, then $\det(A) = \pm 1$.
- (i) If W_1 and W_2 are subspaces of a vector space V , then $W_1 \cup W_2$ is also a subspace of V .
- (j) Let A and B be $m \times n$ and $n \times m$ matrices respectively. Then AB and BA are well defined and $\text{rank}(AB) = \text{rank}(BA)$.
- (k) Let A be 3×3 matrix. If A has only two distinct eigenvalues λ_1 and λ_2 , then A is not diagonalizable.
- (l) Suppose $A_{n \times n}$ is similar to a diagonal matrix $D_{n \times n} = [d_{ij}]_{1 \leq i, j \leq n}$. Namely, $A = PDP^{-1}$ for some nonsingular matrix $P_{n \times n}$. Then $\sum_{k=1}^{\infty} A^k$ exists if $|d_{ii}| < 1$ for all $i = 1, 2, \dots, n$.

考 試 科 目	基礎數學	系 所 別	統計學系	考 試 時 間	2 月 9 日 (三) 第一節
2. (12pts). Please sketch the graph of the polynomial $f(x) = 4x^5 + x^3 - 2x + 1$ and find the integer a such that the only real root of $4x^5 + x^3 - 2x + 1 = 0$ is in $(a, a + 1)$. Show your work. 3. (10pts). Please use appropriate power series to show $\ln 2 = \sum_{k=1}^{\infty} \frac{1}{k2^k}$. 4. (10pts). Please evaluate $\iint_D \frac{(x+2y)\sqrt{x-y+3}}{x+2y+1} dA$ by making an appropriate change of variables, where D is the parallelogram enclosed by the lines $x + 2y = 0, x + 2y = 4, x - y = 1$, and $x - y = 6$. Show your work. 5. (12pts). Let W be the subspace of $\mathcal{R}^4$ spanned by the vectors $\mathbf{w}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{w}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \\ 2 \end{bmatrix}$, and $\mathbf{w}_3 = \begin{bmatrix} -1 \\ 2 \\ 3 \\ 0 \end{bmatrix}$. Please find the (orthogonal) projection of $\mathbf{v} = \begin{bmatrix} 4 \\ 1 \\ 0 \\ -1 \end{bmatrix}$ on W. Show your work. 6. (10pts). Consider the vector space P_2 of all polynomials of degree ≤ 2 together with the zero polynomial. Note $S = \{p_1(t) = t^2 + t + 1, p_2(t) = t + 1, p_3(t) = 1\}$ is a basis for P_2. Let $L : P_2 \rightarrow P_2$ be the linear transformation whose matrix with respect to S is $\begin{bmatrix} 0 & 0 & 0 \\ 2 & 0 & 0 \\ -1 & 1 & 0 \end{bmatrix}$. Please find $L(at^2 + bt + c)$. Show your work. 7. (10pts). Records about the migration patterns of a species of bird from year to year among three habitats A, B and C show that of the birds beginning of the year in habitat A, 20% migrate to habitat B, and 50% migrate to habitat C. Of the birds beginning of the year in habitat B, 10% migrate to habitat A, and 10% migrate to habitat C. Of the birds beginning of the year in habitat C, 40% migrate to habitat A, and 40% migrate to habitat B. In the long wrong, what percentage of the birds will live in each of the habitats? Show your work.					
備 註	一、作答於試題上者, 不予計分。 二、試題請隨卷繳交。				

考 試 科 目	數理統計學	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第二節
1. (23%) Two machines of the same type, A and B, always operate under the same environment, but an extra filter had installed on A to reduce the pollutant. Let Y_1 be the amount of pollutant emitted by A, while Y_2 be the pollutant amount emitted by B. The joint pdf is $f(y_1, y_2) = \begin{cases} 1/2, & 0 \leq y_1 \leq 1, 0 \leq y_2 \leq 4, 4y_1 \leq y_2, \\ 0, & \text{otherwise.} \end{cases}$ (1) Find the probability that Y_2 / Y_1 is at least 5. (5%) (2) Find the condition pdf of Y_1 given $Y_2 = y_2$, and $V(Y_1 y_2)$. (8%) (3) The reduction of pollutant amount due to the filter is $R = Y_2 - Y_1$. Find the probability density function of R and its mean. (10%) 2. (22%) Let $X_1, \dots, X_n$ be a random sample from $f(x; \theta) = \frac{1}{\theta}, x \leq \frac{\theta}{2}; \theta > 0$. (1) Find the complete and sufficient statistic for θ. (10%) (2) Find the MVUE (minimum variance unbiased estimator) of $\theta^{-1} + \theta$. (12%) 3. (15%) Given that $N = n$, the conditional pdf of X is $f(x n) = \frac{x^{n-1}e^{-x}}{\Gamma(n)}, x > 0$. Moreover, $P(N=n) = \frac{\mu^n e^{-\mu}}{n!}, n = 0, 1, 2, \dots$ (1) Find the unconditional mean and variance of X, i.e. $E(X)$ and $V(X)$. (5%) (2) Find the limiting distribution for $\frac{X - E(X)}{\sqrt{V(X)}}$ as $\mu \rightarrow \infty$. (10%) 4. (40%) Let $X_1, \dots, X_n$ be a random sample from $f(x; \theta) = \lambda x^{-(\lambda+1)}, x > 1; \lambda > 0$. (1) Find the MLE (maximum likelihood estimator) of λ. (5%) (2) Find the Cramer-Rao lower bound for the variance of unbiased estimators of $P(X > a)$, where a is a constant greater than 1. (7%) (3) Find the UMP size α test of $H_0: \lambda \leq \lambda_0$ versus $H_1: \lambda > \lambda_0$. (10%) (4) Find the $100(1-\alpha)\%$ confidence interval for λ. (6%) (5) Let $Y_1, \dots, Y_m$ be another random sample from $f(y; \theta) = \theta y^{-(\theta+1)}, y > 1; \theta > 0$. Derive the likelihood ratio test of $H_0: \lambda = \theta$ versus $H_1: \lambda \neq \theta$. (12%)					
備	註	一、作答於試題上者, 不予計分。 二、試題請隨卷繳交。			

考 試 科 目	統計方法	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第 四 節
---------	------	-------	------	---------	------------------

1. (12%) A Center of Sleep Disorder conducted a survey to determine whether the hours of sleep per night are independent of age. A random sample of 500 individuals was asked two questions: (i) hours of sleep per night (with four options: fewer than 6, 6 to 6.9, 7 to 7.9, and 8 or more); (ii) age (younger than 30, and 30 or older). The data are given in the following table.

Hours of sleep	Age	
	Younger than 30	30 or older
fewer than 6	60	90
6 to 6.9	70	70
7 to 7.9	60	60
8 or more	50	40

- (a) Using a 0.05 significance level, conduct a hypothesis test to determine whether the hours of sleep per night are independent of age. Please complete the hypothesis test. (10%)
- (b) What is your interpretation about the result in (a)? (2%)
2. (20%) For testing $H_0: \mu \leq 22$ v.s. $H_a: \mu > 22$, a random sample of 256 individuals is selected from the population. The population variance is 400. The power is 0.2005, denoted by $1 - \beta_{23.4}$, when the actual population mean is 23.4.
- (a) What is the significance level adopted in the test? (12%)
- (b) In such a hypothesis testing with a sample of 256 individuals, is it possible that the probability of type II error $\beta_{23.4}$ and the significance level decrease simultaneously? Please give your reason. (8%)
3. (35%) A realty company sells apartments in the Midwest of the United States. Since their customers are interested in the fee they have to pay for heating in winter, a research group in the company collected data and tried to build regression models to predict the heating cost (abbreviated to Heat.cost; denoted by Y). There were three independent variables: the mean temperature in winter (Temp, X_1), the centimeters of insulation in the attic (Insu, X_2), and the age (years) of the heater (Age, X_3). The research group considered the following regression analyses:
- Model 1: $Y = \alpha + \beta_1 X_1 + \eta$,
- Model 2: $Y = \alpha + \beta_3 X_3 + \delta$,
- Model 3: $Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \varepsilon$,

考 試 科 目	統計方法	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第 四 節
---------	------	-------	------	---------	------------------

where η , δ , and ε are error terms. They collected data from 20 of their former customers for data analysis. First of all, the Model 1 is fitted based on the 10 elements given in the following table, and please answer Questions (a) and (b).

Customer	Y	X_1
1	250	-5
2	360	-9
3	165	-5
4	43	9
5	92	11
6	200	-8
7	355	-19
8	290	-21
9	230	-13
10	120	6

(a) Find the least squares estimates of α and β_1 under Model 1. (12%)

(b) Compute the coefficient of determination, R^2 , for Model 1. (12%)

An output from a computer software for fitting Model 3 to the data is displayed in the followings. Using the output to answer Questions (c) and (d).

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	228.833	40.547	5.644	3.66e-05 ***
Temp	-8.348	1.361	-6.134	1.44e-05 ***
Insu	-6.098	1.872	-3.258	0.00494 **
Age	5.751	3.937	1.461	0.16349

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 49.98 on 16 degrees of freedom

Multiple R-squared: 0.8123, Adjusted R-squared: 0.7771

F-statistic: 23.08 on 3 and 16 DF, p-value: 4.695e-06

The output for fitting Model 2 to the data is displayed in the followings.

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	86.184	48.656	1.771	0.0934 .
Age	17.009	6.303	2.699	0.0147 *

考 試 科 目	統計方法	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第 四 節
---------	------	-------	------	---------	------------------

(c) What is your interpretation about the result of the F-test for Model 3 ? (5%)

(d) Based on the fittings, the β_3 under Model 2 is significant while the β_3 under Model 3 is not, at a 0.05 significance level. Comment on the phenomenon (Is there any contradiction, or ...?). (6%)

4. (10%) Consider the hypothesis test that $H_0: \mu \geq \mu_0$ v.s. $H_a: \mu < \mu_0$, where μ_0 is a fixed constant. The population variance is known as σ^2 . Suppose that the significance level is set at α , and suppose further that the probability of type II error is β when the population mean is μ_{0a} , where μ_{0a} is a constant ^{smaller} than μ_0 . Define z_α and z_β as numbers satisfying $P(Z \geq z_\alpha) = \alpha$ and $P(Z \geq z_\beta) = \beta$. Derive that the sample size should be $\frac{(z_\alpha + z_\beta)^2 \cdot \sigma^2}{(\mu_0 - \mu_{0a})^2}$ (and then round up).

5. (23%) A research study is conducted to compare the kilometers per liter of unleaded regular, mid-grade, and super premium gasolines. There are differences in the performances of different automobiles. Seven different automobiles (numbered in 1 to 7) are sampled randomly and tested. The experimental results, in kilometers per liter, are given in the following table.


Automobile	Gasolines		
	Regular	Mid-grade	Super Premium
1	26	26	29
2	28	25	28
3	26	28	30
4	29	28	30
5	31	30	35
6	28	29	33
7	28	30	32

- (a) To answer the research question, an ANOVA is performed. What are the null and alternative hypotheses for the ANOVA? (2%)
- (b) Set up the ANOVA table. (17%)
- (c) Using a 0.05 significance level, what is the conclusion? What is your interpretation about the result? (4%)

考 試 科 目	統計方法	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第 四 節
---------	------	-------	------	---------	------------------

Table of the standard Normal Distribution (its cumulative probabilities)

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641




考 試 科 目	統計方法	系 所 別	統計學系	考 試 時 間	2 月 9 日(三) 第 四 節
---------	------	-------	------	---------	------------------

Table of the F Distribution ($\alpha = 0.05$)

		Degrees of Freedom for the Numerator															
		1	2	3	4	5	6	7	8	9	10	12	15	20	24	30	40
Degrees of Freedom for the Denominator	1	161	200	216	225	230	234	237	239	241	242	244	246	248	249	250	251
	2	18.5	19.0	19.2	19.2	19.3	19.3	19.4	19.4	19.4	19.4	19.4	19.4	19.4	19.5	19.5	19.5
	3	10.1	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.66	8.64	8.62	8.59
	4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.80	5.77	5.75	5.72
	5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.62	4.56	4.53	4.50	4.46
	6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.94	3.87	3.84	3.81	3.77
	7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.51	3.44	3.41	3.38	3.34
	8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.22	3.15	3.12	3.08	3.04
	9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.01	2.94	2.90	2.86	2.83
	10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.85	2.77	2.74	2.70	2.66
	11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.72	2.65	2.61	2.57	2.53
	12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.62	2.54	2.51	2.47	2.43
	13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.53	2.46	2.42	2.38	2.34
	14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.46	2.39	2.35	2.31	2.27
	15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.40	2.33	2.29	2.25	2.20
	16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.42	2.35	2.28	2.24	2.19	2.15
	17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.38	2.31	2.23	2.19	2.15	2.10
	18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.34	2.27	2.19	2.15	2.11	2.06
	19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.31	2.23	2.16	2.11	2.07	2.03
	20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.28	2.20	2.12	2.08	2.04	1.99

Table of the Chi-square Distribution



Degrees of Freedom	Area in Upper Tail									
	.995	.99	.975	.95	.90	.10	.05	.025	.01	.005
1	.000	.000	.001	.004	.016	2.706	3.841	5.024	6.635	7.879
2	.010	.020	.051	.103	.211	4.605	5.991	7.378	9.210	10.597
3	.072	.115	.216	.352	.584	6.251	7.815	9.348	11.345	12.838
4	.207	.297	.484	.711	1.064	7.779	9.488	11.143	13.277	14.860
5	.412	.554	.831	1.145	1.610	9.236	11.070	12.832	15.086	16.750
6	.676	.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.647	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.041	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801

備

註

- 一、作答於試題上者，不予計分。
二、試題請隨卷繳交。